## Families of Fungicides for Turfgrass J. W. Rimelspach and T. E. Hicks, The Ohio State University, Department of Plant Pathology

turfdisease.osu.edu

Common Name	FRAC Code <sup>2</sup>	Trade Names <sup>1</sup>	Mode of Action	Uptake and/or Mobility	Concern Over Resistance	Comments		
		C	hemica	I Family: Dithic	ocarbamates			
Mancozeb	M3	Fore, Mancozeb, Dithane T/O, Protect T/O	general	contact (no uptake into the tissue)	low	These types of fungicides have broad-spectrum control properties and are used as protectants.  Early development of these started in the 1930s.		
Thiram	M3	Spotrete, Defiant, Thiram	Family #					
	Π	Cnemical	Family:	Nitriies/Benzo	onitriles/Chloror			
Chlorothalonil	M5	Daconil, Manicure, Pegasus, Echo, QP Chlorothalonil	general	contact (no uptake into the tissue)	low	Introduced in the late 1960s and now used extensively on many crops worldwide. This chemistry can provide excellent protection for many infectious diseases, but cannot suppress existing infections. Proper application technique is a must. There are no reports of resistance.		
			Chemic	al Family: Ben	zimidazoles			
Thiophanatemethyl	1	3336, thiophanate methyl T methyl Pro, T-Storm	specific	systemic (upward)	high	This family of fungicides became available in the late 1960s and ushered in the era of systemic fungicides. The development of resistance to the benzimidazoles is a serious problem.		
	Chemical Family: Dicarboximides							
Iprodione	2	Chipco 26GT, Raven Iprodione Pro, 18 Plus, QP Ipro	specific	local penetrant	moderate to high (not persistent)	The dicarboximides were developed in the mid-1970s. These fungicides have broad-spectrum activity.		
Vinclozolin	2	Chaminal Family	Charall		Domosthy does Ind	sibitoro (DMI)		
Egnarimol	2	Chemical Family:  Rubigan <sup>3</sup>	Steroi I	mmonors (SI)/L	Jenneunylase int	IIDITOIS (DIVII)		
Fenarimol  Myclobutanil	3	Eagle, QP Myclobutanil	-					
Triademefon	3	Bayleton, Accost	-					
Propiconazole	3	Banner MAXX, Spectator, ProPensity, Kestrol, ProPimax, QP Propiconazole	specific	systemic (upward)	high	This group of fungicides was introduced in the late 1970s and has broad-spectrum activity. At times, referred to as the SIs or DMIs. The development of resistance to this family of fungicides is a problem.		
Triticonazole	3	Trinity, Triton <sup>3</sup>	1					
Metconazole	3	Tourney	]					
Tebuconazole	3	Torque, Mirage, Stressgard						
Mefentrifluconazole	3	Maxtima						
Flutriafol	3	Rayora						
	l .		T T		nate Dehydroge	nase Inhibitors (SDHI)		
Flutolanil	7	ProStar, Pedigree	specific	systemic (upward)	low	-		
Boscalid Fluxapyroxad	7	Emerald  Xzemplar	specific specific	systemic (upward) systemic (upward)	moderate moderate	The products listed have similar target sites; however,		
Penthiopyrad	7	Velista	specific	systemic (upward)	moderate	they are typically used to manage different diseases.  Newer materials are active on a broad range of turfgrass		
Isofetamid	7	Kabuto	specific	systemic (upward)	moderate	diseases.		
Pydiflumetofen	7	Posterity	specific	systemic (upward)	moderate	-		
		C	hemica	l Family: Strok	oilurins (Qol)			
Azoxystrobin	11	Heritage	specific	systemic (upward)	high			
Trifloxystrobin	11	Compass	specific	local penetrant	high	Azoxystrobin was introduced in 1997 and the chemical		
Pyraclostrobin	11	Insignia	specific	local penetrant	high	structures were produced by various naturally occurring, wood-decaying fungi. Strobilurins are broad spectrum		
Fluoxastrobin	11	Disarm, Fame	specific	systemic (upward)	high	disease management tools.		
Mandestrobin	11	Pinpoint	specific	systemic (upward)	high			
			Chemi	cal Family: Phe	enylpyrrole			
Fludioxonil	12	Medallion	specific	local penetrant	low to moderate	Enters the turf plant and is translaminar; it moves from one leaf surface to the other side of leaf.  Does not move in the xylem.		
		Chei	nical Fa	amily: Aromati	c Hydrocarbons			
PCNB or Quintozene	14	Terraclor, Turfcide, Revere, FFII, PCNB, Defend, Engage	general	contact (no uptake into the tissue)	low	PCNB is usually considered to be a protectant but may be locally systemic. Considerable label changes are occurring at this time.		
			Che	mical Family:	Polyoxin			
Polyoxin D zinc salt	19	Endorse, Affirm	specific	local penetrant	moderate	The fungicide enters the plant tissue and accumulates in the waxy cuticle and has translaminar movement. Polyoxin D can suppress existing fungal infections.		
			Chemi	cal Family: Pyr	ridinamine			
Fluazinam	29	Secure	general	contact (no uptake into the tissue)	low	A new, contact, multi-site, broad spectrum fungicide introduced into other crops in the 1990s and turf in 2012.		

(continued on back for Pythium/Oomycete Materials and Combination Fungicide Products)



		The follow	<i>i</i> ing are	used for Pythium and	water molds	
Common Name	FRAC Code <sup>2</sup>	Trade Names <sup>1</sup>	Mode of Action	Uptake and/or Mobility	Concern Over Resistance	Comments
	Few diseases besides those					
Mefenoxam	4	Subdue MAXX, QP Mefenoxam Apron (seed treatment)	specific	systemic (upward)	high	caused by Pythium species or closely related water molds (Oomycetes), like yellow tuft, are controlled. Azoxystrobin (Heritage)
	and Pyraclostrobin (Insignia) have unique activity against both					
Azoxystrobin	11	Heritage	specific	systemic (upward)	moderate to high	Pythium species (Oomycetes)
Pyraclostrobin	11	Insignia	specific	local penetrant	high	and true fungi. Fosetyl-aluminum is a true systemic exhibiting both
Chemical Family: Aromatic Hydrocarbons						upward and downward movement in plants. It is also unique in that it
Chloroneb	14	Teremec SP	general	contact (local penetrant)	low	moves in the phloem (symplastic transport) as compared to all
Ethazole (Etridiazole)	14	Koban, Terrazole, Truban	general	contact	low	other systemic fungicides that are transported in the xylem (apoplastic transport).
		Chemical Fa	amily: C	yanoimidazole		- transporty.
Cyazofamid	21	Segway	specific	local penetrant	moderate to high	
		Chemical	Family	: Carbamate		
Propamocarb	28	Banol	not well known	systemic (upward)	low	
Chemical Family: Phosphonate						
Fosetyl-aluminum	P7	Prodigy, Chipco Signature, QP Fosetyl- Al	not well known	systemic (up and down)	low	
phosphite (salts of phosphorous acid)	P7	Magellan, Biophos, Resyst, Alude, Vital, Kphite, Fiata Stressgard, Appear	general	systemic (up and down)	low	
Chemical Family: Benzamide and Carbamate						
Fluopicolide	43+28	Stellar (combined with propamocarb)	general	systemic (upward)	low	

<sup>1</sup>Product list by trade name may not be all inclusive. No endorsement is intended for products mentioned or is criticism meant for products not mentioned.

<sup>&</sup>lt;sup>3</sup>Product no longer manufactured for turfgrass use.

Combination Fungicide Products					
Product Name (Trade Names)	FRAC Codes	Active Ingredients by Common Names			
Armada	(3 + 11)	triadimefon + trifloxystrobin			
Briskway	(11 + 3)	azoxystrobin + difenoconazole*			
Civitas One	_	synthetic isoparaffin + other ingredients			
Concert	(3 + M5)	propiconazole + chlorothalonil			
Consan	_	dimethyl benzyl + dimethyl ethylbenzyl, ammonium chlorides			
ConSyst, Spectro, Peregrine	(1 + M5)	thiophanate-methyl + chlorothalonil			
Daconil ACTION	(M5 + P1)	chlorothalonil + acibenzolar-S-methyl			
Enclave	(M5 +2 + 1 + 3)	chlorothalonil + iprodione + T-methyl + tebuconazole			
Exteris Stressgard	(7 + 11)	fluropyram* + trifloxystrobin + Stressgard			
Fame +C	(11 + M5)	fluoxastrobin + chlorothalonil			
Fame +T	(11 + 3)	fluoxastrobin + tebuconazole			
Headway	(11 + 3)	azoxystrobin + propiconazole			
Honor	(11 + 7)	pyraclostrobin + boscolid			
Instrata	(3 + M5 + 12)	propiconazole + chlorothalonil + fludioxonil			
Interface Stressgard	(2 + 11)	iprodione + trifloxystrobin + Stressgard			
Junction	(M1 + M3)	copper hydroxide + mancozeb			
Lexicon	(7 + 11)	fluxapyroxad + pyraclostrobin			
MANhandle	(3 + M3)	myclobutanil + mancozeb			
Navicon	(3 + 11)	mefentrifluconazole + pyraclostrobin			
Pillar	(11 + 3)	pyraclostrobin + triticonazole			
Premion	(14 + 3)	PCNB + tebuconazole			
Prostar Plus	(3 + 7)	triadimefon + flutolanil			
Renown	(11 + M5)	azoxystrobin + chlorothalonil			
Stellar	(43 + 28)	fluopiolide + propamocarb hydrochloride			
Systar	(1 + 7)	thiophanate-methyl + flutolanil			
Tartan Stressgard	(3 + 11)	triadimefon + trifloxystrobin + Stressgard			
Tekken	(7 + 3)	isofetamid + tebuconazole			
Traction	(29 + 3)	fluazinam + tebuconazole			
26/36 Fungicide, Lesco TwoSome	(2 + 1)	iprodione + thiophanate-methyl			
*This is NOT sold as a single material for turfgrass.					

Biocontrol Agents				
Product Name (by Trade Names)	Active Ingredients by Common Names			
EcoGuard	Bacillus licheniformis			
Companion	Bacillus subtillis GB03			
Rhapsody	QST 713 strain of <i>Bacillus subtilis</i>			
TurfShield	Trichoderma harzianum Rifai strain T-22 + Trichoderma virens strain G-41			

FRAC (Fungicide Resistance Action Committee)

FRAC is a Specialist Technical Group of CropLife International

FRAC Code: Numbers and letters are used to distinguish the fungicide groups according to their cross resistance behavior. The numbers were assigned according to the time of product introduction to the market. The letters refer to: P = host defense inducers, M = multi-site inhibitors, and U = unknown mode of action and unknown resistance risk. For more information, go to **frac.info/home**.

For additional information, refer to our website: turfdisease.osu.edu.

<sup>&</sup>lt;sup>2</sup>FRAC codes indicate the biochemical target site of action, according to the Fungicide Resistance Action Committee. M3 and M5 indicate multi-site inhibitor, with no significant risk of resistance.